Comprehensive evaluation of the cytotoxicity of CdSe/ZnS quantum dots in *Phanerochaete chrysosporium* by cellular uptake and oxidative stress

Liang Hu^{a,b}, Jia Wan^{a,b}, Guangming Zeng^{a,b,*}, Anwei Chen^{c,*}, Guiqiu Chen^{a,b}, Zhenzhen Huang^{a,b}, Kai He^{a,b}, Min Cheng^{a,b}, Chengyun Zhou^{a,b}, Weiping Xiong^{a,b}, Cui Lai^{a,b}, Piao Xu^{a,b}

^a College of Environmental Science and Engineering, Hunan Oniversity, Changsha, Hunan

410082, P.R. China

^b Key Laboratory of Environmental Biology and Pollution Control (Hunan University), Ministry of Education, Changsha, Hunan 410082, P.R. Chine

^c College of Resources and Environment, Hunan Agricultural University, Changsha, Hunan

410128, P.R. China

*Corresponding author: Address: College of Environmental Science and Engineering, Hunan University, Changsha 410082, P.R. China. Tel.: +86 731 88822829; fax: +86 731 88823701.

E-mail addresses: zgming@hnu.edu.cn.

Abstract

The growing potential of quantum dots (QDs) in biological and biomedical applications has raised considerable concerns due to their toxicological impact. Consequently, it is urgent to elucidate the underlying toxicity mechanism of QDs. In this work, we comprehensively investigated the cellular uptake and induced physiological responses of four CdSe/ZnS QDs (COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625) in Phanerechaete chrysosporium (P. chrysosporium) through using inductively coupled plasma optical emission spectrometer, confocal laser scanning microscopy and the analyses of malondialdehyde content, superoxide level, superoxide dismutase activity, catalase activity and glutathione level. The results howed that four CdSe/ZnS QDs accumulated largely in the hyphae and caused oxidative stress to P. chrysosporium in the tested concentration range (10-80 nM). Furthermore, the cellular uptake and cytotoxicity were related to the physicochemical properties of QDs, such as particle size and surface charges. Negative charged CdSe/ZnS QDs with small size could be *P. chrysosporium* than large ones, thus small size CdSe/ZnS more easily more cytotoxic to P. chrysosporium. On the other hand, small negative ODs were charged CdSe/ZnS QDs resulted in greater cytotoxicity than large negative charged CdSe/ZnS QDs. The obtained results offer valuable information for revealing the toxicity mechanism of QDs in living cells.

Keywords

CdSe/ZnS quantum dots; Phanerochaete chrysosporium; Cytotoxicity; Cellular

uptake; Oxidative stress; Inductively coupled plasma optical emission spectrometer

1. Introduction

Quantum dots (QDs) are generally made up of atoms from IIIA-VA or IIB-VIA elements in the chemical periodic table.¹ A semiconductor core (e.g., CdS, CdSe, and CdTe) which can be encapsulated in a shell (e.g., ZnS) to enhance both electronic and optical properties and reduce core metal leaching is the main structure of a typical QD.^{2,3} QDs are also among the most promising fluorescent hanoparticles for biological and biomedical applications because of her unique photophysical properties, such as high brightness, tunable broad excitation coupled with narrow emission spectra, and excellent photostability. With the recent development of bioconjugate technique and surface modification, QDs have been extensively applied to *in vitro* and *in vivo* imaging, virus and cell tracing, cellular proteins labelling, targeted drug delivery, and cancer therapy.⁹⁻¹¹ However, the comprehensive toxicity evaluation of QDr is of especial significance for their clinical medicine due to their heavy metal components and nanosize effects.

Although the potential toxicity of QDs has remained large controversies in biomedical applications and challenges for clinical studies, many toxicological studies on QDs have been implemented for this purpose.^{12,13} It has been verified that the cytotoxicity of QDs would lead to cell growth inhibition, mitochondrial dysfunction, DNA damage, and apoptosis.¹⁴⁻¹⁶ The physicochemical properties of QDs, including core composition, size, surface charge, and functionalization, would influence their

cytotoxicity to a great extent.¹⁷ For example, Li et al.¹⁸ reported that CdSe QDs were more cytotoxic than CdSe/ZnS QDs with the same size and functionalization, indicating that a cadmium-induced cytotoxic response could be decreased by encapsulating the core with a shell. The finding was further supported by Su et al.,¹⁹ who showed that CdTe/ZnS QDs were less cytotoxic as compared to CdTe QDs. In the same way, Domingos et al.²⁰ reported that CdTe QDs capped with surfactant were more cytotoxic than CdTe/ZnS QDs coated with polyethylene giveol (PEG) on *Chlamydomonas reinhardtii* cells. In regard to size effect, Soenen et al.²¹ found that the cellular uptake and subcellular distribution of CdTe QDs changed dramatically as a result of QDs size. There are also some studies which showed that surface charge of QD could affect their cytotoxicity,^{15,22} with nostive charged QDs being more cytotoxic than negative charged QDs.¹ withionally, QDs functionalization could alleviate the induced cytotoxicity as well.^{23,24}

To further explore and understand the cytotoxicity of QDs, many research groups have focused their attention on the mechanism study.^{12,13,25} It has been widely deemed that the release of toxic cadmium ion (Cd²⁺) and the generation of reactive oxygen species (**POS**) are the main reasons that responsible for QDs cytotoxicity.^{14,26,27} Since Cd²⁺ can be released through the oxidation of QDs and bind to the sulfhydryl groups in many intracellular proteins, it may result in the functionality reduction of various subcellular organelles.^{20,28} QDs-induced ROS has been verified to cause metabolic functions loss, DNA nicking and break, and apoptosis.^{9,10,29} However, it is difficult to thoroughly understand the potential cytotoxicity mechanism of QDs since the cytotoxicity caused by QDs is extremely complicated. Few consensuses can be reached. Although each of above studies has provided valuable information on the influence of a given property of QDs (i.e., core composition, size, surface charge, and functionalization), a comparative analysis is necessary to determine the extent to which each of these properties causes cellular responses in isolation or in combination. It is still unclear which specific properties of QDs lead the induced physiological responses.

In view of this, four types of CdSe/ZnS QDs in this study were employed as different types of pollutants to investigate their cellular uptake and induced physiological responses in Phanerochaete chrysosporium (*P. chrysosporium*), which has been widely used in treating wastewater containing toxic organic pollutants and heavy metals because of its excellent ability o degrade organic pollutants and absorb heavy metals.³⁰⁻³² As P. chryso. portum was sensitive to xenobiotics and it could surrounding environments, the P. chrysosporium quickly respond to the was used as the target organism to explore the cytotoxicity of QDs. Moreover, the physiological responses of P. chrysosporium under QDs exposure are still limited. the effects of incubation concentration and exposure time of QDs on the Particular cellular uptake and oxidative stress in P. chrysosporium have not been reported in the literature yet. Consequently, a simultaneous analysis of the cellular uptake and physiological responses of P. chrysosporium is determined to obtain a reliable and comprehensive evaluation on the biosafety of QDs.

2. Materials and methods

2.1. Reagents and instruments

QDs used in this work were purchased from Wuhan Jiayuan Quantum Dots Company (Wuhan, China). The nanoparticles were preserved as 8 μ M QD in 200 μ L borate buffer solutions. Four CdSe/ZnS QDs (COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625) consist of cadmium selenide (CdSe) core and zinc sulfide (ZnS) shell, encapsulated with the uniform amphipathic polymer of PEG coating. Their differences are particle size and surface functional groups, i.e. -NH₂ and -COOH. In this work all reagents must be of analytical reagent grade and were purchased from Sigma (St. Louis, MO, USA). Ultrapure water produced by a Milli-Q system (18.25 M Ω cm⁻¹, Millipore, France) was used throughout the process. To control the blank value, the experiment ware was completely washed by solving m 0% nitric acid (HNO₃) for at least 24 h.

CdSe/ZnS QDs were characterized by photoluminescence (PL), high-resolution transmission electronic microscopy (HRTEM), and dynamic light scatterer (DLS), respectively. Ill optical measurements were carried out at room temperature under ambient air conditions. PL measurements were performed using a HORIBA JOBTN YVON FLUOROMAX-4 spectrofluorimeter. The PL quantum yield (QY) of samples was estimated using Rhodamine 6G (QY = 95%) in ethanol solution as a reference standard, which was freshly prepared to decrease the measurement error.³³ The HRTEM overview images were recorded with a Philips CM 200 electron microscope (JEOL JEM-3010, Japan) operated at 200 kV. DLS analysis (hydrodynamic diameter and zeta (ζ) potential) was carried out using a DynaPro Dynamic Light Scatterer (Malvern Instruments).

2.2. P. chrysosporium culture

P. chrysosporium has been selected as the objective microbe due to its extensive utilization in wastewater decontamination. *P. chrysosporium* BKMF-1767 (ATCC 24725) was obtained from the China Center for Type Culture Collection (Wuhan, China). The fungal spores were prepared by subculturing on potato dextrose agar slants (20.0 g L⁻¹ glucose, 20.0 g L⁻¹ agar, 3.0 g h⁻¹ EH₂PO₄, and 1.5 g L⁻¹ MgSO₄·7H₂O) in sterilization work station, and then these potato dextrose agar slants were placed in humidity incubator at 37°C for 7 days. Fungal spore suspensions were adjusted to 2.0×10^6 CFU mL⁻¹ using a turbidimeter (WGZ-200, Shanghai, China). 3 mL of aqueous suspensions of fungal spores were inoculated into 200 mL Kirk's liquid culture mediumara 37°C for 3 days.³⁴

2.3. Inductively coupled plasma optical emission spectrometer analysis

0.2 **c** of *P. chrysosporium* pellets was seeded into 10 mL centrifuge tubes containing fresh borate buffer solutions with CdSe/ZnS QDs. Then these centrifuge tubes were placed in an orbital shaker (120 rpm) at 37 °C. After 24 h of exposure, *P. chrysosporium* pellets were harvested and washed for three times with ultrapure water, and then the pellets were digested with 4 mL concentrated HNO₃ (67%) for 4 h at 85 °C. The solution was evaporated to remove superfluous acid and diluted with 5 mL HNO₃ (5%) for inductively coupled plasma optical emission spectrometer (ICP-OES) (Optima 5300 DV) analysis. The blank sample was the *P. chrysosporium* pellets without CdSe/ZnS QDs exposure. In this work, the uptake amount of CdSe/ZnS QDs was calculated from Cd or Se measured by ICP-OES.

2.4. P. chrysosporium uptake of CdSe/ZnS QDs

2.4.1. Incubation concentrations

0.2 g of *P. chrysosporium* pellets was seeded into 10 mL centrifuge tubes and fresh borate buffer solutions containing different concentrations of CdSe/ZnS QDs (0, 10, 20, 50, and 80 nM) were added. Then these centrifuge tubes were placed in an orbital shaker (120 rpm) at 37 °C. After 24 h of exposure, *P. chrysosporium* pellets were collected and washed for three tubes with ultrapure water. The uptake of CdSe/ZnS QDs by *P. chrysosporium* was determined as described above.

2.4.2. Incubation time

0.2 g of *P elevyposporium* pellets was seeded into 10 mL centrifuge tubes and fresh bornte buffer solutions containing different concentrations of CdSe/ZnS QDs (0, 10, 20, 50, and 80 nM) were added. Then these centrifuge tubes were placed in an orbital shaker (120 rpm) at 37 °C. After incubation for different durations (0, 3, 6, 9, 15, and 24 h), *P. chrysosporium* pellets were collected and washed for three times with ultrapure water. The uptake of CdSe/ZnS QDs by *P. chrysosporium* was determined as described above.

2.5. Confocal laser scanning microscopy

0.2 g of *P. chrysosporium* pellets was seeded into 10 mL centrifuge tubes and 5 mL 2,7-dichlorodihydrofluorescein-diacetate (H₂DCF-DA) (5 μ M) was added into incubation solution for 2 h. Afterward, the medium was discarded, and the pellets were treated with different concentrations of CdSe/ZnS (0, 10, 20, 50, and 80 nM) for 24 h. Then the pellets were washed with fresh phosphate-buffered saline (PBS) three times for confocal laser scanning microscopy analysis (FV1,000, TM1,310 Japan) equipped with double photon detector.

2.6. Lipid peroxidation and superoxide (O_2) production

Lipid peroxidation was determined by mastering the content of malondialdehyde (MDA), which is generally used as the indicator of lipid peroxidation and free radical production.³⁵ MDA content was nearured by the thiobarbituric acid (TBA) reaction according to the method reported by Aravind and Prasad.³⁶ The harvested *P. chrysosporium* pelles were homogenized in 10 mL centrifuge tube with 10% trichloroacetic acte and centrifuged at 10,000 rpm for 20 min at 4 °C. Then the supernatative was separated and boiled with TBA for 20 min. The heated supernatant was centrifuged at 8,000 rpm for 10 min, and the absorbance was recorded at 532 and 600 nm using a UV-vis spectrophotometer (Model UV-2550, Shimadzu, Japan).

 O_2^- production was measured according to the method reported by Lei et al..³⁷ The harvested *P. chrysosporium* pellets were homogenized in 10 mL centrifuge tube with 50 mM PBS (pH 7.8) and centrifuged at 10,000 rpm for 15 min at 4 °C. Then 1

mL of the supernatant was mixed with 0.9 mL of 50 mM PBS (pH 7.8) and 0.1 mL of 10 mM hydroxylamine hydrochloride. The reaction system was placed in water bath at 25 °C for 20 min before adding 1 mL of 7 mM α -naphthylamine and 1 mL of 17 mM p-aminobenzenesulfonic acid. Afterward, the absorbance of above mixture was recorded at 530 nm.

2.7. Antioxidant enzymes analysis

Superoxide dismutase (SOD) (convert superoxide to hydrogen peroxide) activity was determined according to the method described in our previous work.³⁵ *P. chrysosporium* pellets were collected by centrifugation and homogenized in 0.05 M PBS (pH 7.8). Then the homogenate was centrifuged ab10,000 rpm for 15 min at 4 °C, and the supernatant was separated for exvine assay. SOD activity was measured by monitoring the inhibition of photochemical reduction of nitroblue tetrazolium chloride (NBT) in a reaction system containing 2.25 mM NBT, 200 mM methionine, 1 M Na₂CO₃, 60 mM riboflavin, 9 nM EDTA, and 0.1 M PBS (pH 7.8). The absorbance was recorded at 560 nfn.

Catalase (CAT) (convert hydrogen peroxide to oxygen and water) activity was assayed in a 3 mL reaction system containing 20 mM H₂O₂, 50 µL enzymatic extract, and 50 mM PBS (pH 7.8), according to the method described by Cavalcanti et al..³⁸ CAT activity was evaluated by measuring the decrease rate in absorbance at 240 nm (molar extinction coefficient $\varepsilon = 36.6 \text{ mM}^{-1} \text{ cm}^{-1}$).

2.8. Measurement of glutathione levels

Glutathione levels were measured using 5,5-V-dithiobis-(2-nitrobenzoic acid)-glutathione disulfide according to the method reported by Rehman and Anjum.³⁹ The harvested *P. chrysosporium* pellets were rinsed with PBS (0.1 M, pH 7.0) plus 0.5 mM EDTA and sonicated in 10 mL centrifuge tube with ice-water bath. The samples were sonicated for 5 s in intervals of 8 s with a total time of 5 min. The sonication was carried out with the output power of 400 W for each sample. The suspension was centrifuged at 15,000 rpm for 10 min at 4 °C, and the suspension was used for measuring the concentration of glutathione. Reducel glutathione (GSH) was determined by adding 2.0 mL PBS to 0.5 mL of an above supernatant, followed by adding 0.5 mL of 3 mM 5-dithio-bis-(2-mecoarzoic acid). After 5 min reaction, the absorbance was recorded at 412 nm.

3. Results and discussion

3.1. The physicochemical characteristics of CdSe/ZnS QDs

In an attempt to explore the individual and/or collective roles of various physicochemical properties of CdSe/ZnS QDs, a range of CdSe/ZnS QDs size, surface charge, and functional group was investigated. Specifically, four types of CdSe/ZnS QDs with maximum luminescent wavelengths of 525 nm, 625 nm, 525 nm, and 625 nm were used in this experiment. With an excitation wavelength of 380 nm, COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625

present fluorescence emission maximum at 524 nm, 620 nm, 522 nm, and 624 nm (Table 1), respectively, indicating that four types CdSe/ZnS QDs are nearly monodisperse and homogeneous. The physical diameters of COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625 determined by HRTEM are 5.2 ± 0.3 , 10.4 ± 1.1 , 5.4 ± 0.5 , and 10.3 ± 0.9 nm (Fig. 1), respectively, which are smaller than those $(22.3 \pm 5.4, 30.1 \pm 3.6, 17.6 \pm 3.5, \text{ and } 27.8 \pm 2.5 \text{ nm})$ determined by DLS. The deviation of diameter measured by HRTEM and DLS is attributed to different surface states of nanoparticles under the tested conditions.^{40,41} Since QDs samples are directly measured in aqueous phase for **D**LS analysis while the solution must be strictly evaporated in HRTEN characterization. The zeta (ζ) potentials of COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH2 CdSe/ZnS 525, and 6.7 ± 0.6 , and -9.6 ± 0.7 mV, NH₂ CdSe/ZnS 625 are -12.9 ± 1.1 , 1.4, respectively. Thus, COOH CdSe/InSQDs possess larger negative charges than that of NH2 CdSe/ZnS QDs. In addition, the amount of Cd and Se atoms in a single QD nanoparticle ($N_{Cd/QD}$ and $N_{Se/QD}$) was determined by ICP-OES (Table 1).

Tabla	1
Table	L

The physicoc	hemical	properties	of four	CdSe/ZnS	QDs.
--------------	---------	------------	---------	----------	------

QDs	Function group	$\lambda_{emission}$ (nm)	Physical diameter (nm)	Hydrodynamic diameter (nm)	Zeta potential (mV)	N _{Cd/QD}	$N_{\text{Se/QD}}$
COOH-QDs 525	СООН	524	5.2 ± 0.3	22.3 ± 5.4	-12.9 ± 1.1	165 ± 4	28 ± 5
COOH-QDs 625	СООН	620	10.4 ± 1.1	30.1 ± 3.6	-15.2 ± 1.4	1240 ± 36	359 ± 24
NH ₂ -QDs 525	NH_2	522	5.4 ± 0.5	17.6 ± 3.5	-6.7 ± 0.6	174 ± 7	32 ± 3
NH ₂ -QDs 625	NH ₂	624	10.3 ± 0.9	27.8 ± 2.5	-9.6 ± 0.7	1263 ± 45	343 ± 33



Fig. 1. High-resolution TEM images of COOH CdSe/ZnS 525 (a), COOH CdSe/ZnS 625 (b), NH₂ CdSe/ZnS 525 (c), and NH₂ CdSe/ZnS 625 (d), respectively.

3.2. P. chrysosporium uptake of CdSe/ZnS QDs

3.2.1. Incubation concentrations

The cellular uptake process plays a key role in the function of QDs in biomedical application and their health risks. In this study, chrysosporium pellets were incubated with different concentrations of SOOP CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625 (10, 20, 50, and 80 nM for each QD) for 24 h, and the cellular uptake of CdSe/ZnS QDs was determined by ICP-OES. As shown in Fig. 2a, the uptake of P. chrysosporium increased gradually with the increase of incubation concentration for all four types of CdSe/ZnS QDs. With the ertration of QDs, the amount of CdSe/ZnS QDs in cell membrane increasing increased, leading to the increasing uptake of CdSe/ZnS QDs by P. chrysosporium. However, the cellular uptake of different CdSe/ZnS QDs was distinctly different under the identical conditions. For example, the cellular uptake of NH₂ CdSe/ZnS 525 reached 292.3 pmol g⁻¹ biomass when the incubation concentration was 80 nM, which was greater than those of NH₂ CdSe/ZnS 625 (166.5 pmol g⁻¹ biomass). COOH CdSe/ZnS 525 (101.2 pmol g⁻¹ biomass), and COOH CdSe/ZnS 625 (67.5 pmol g⁻¹

biomass). Among the four types of CdSe/ZnS QDs, the uptake of P. chrysosporium decreased as follows: NH₂ CdSe/ZnS 525 > NH₂ CdSe/ZnS 625 > COOH CdSe/ZnS 525 > COOH CdSe/ZnS 625, which suggested greater uptake of amino-QDs by P. chrysosporium. The result may attribute to that amino-QDs possess fewer surface negative charges (- 6.7 ± 0.6 and - 9.6 ± 0.7 mV) than that (- 12.9 ± 1.1 and - 15.2 ± 1.4 mV) of carboxyl-QDs. Due to the electrostatic repulsion caused by the negative charges on cell membrane, the larger surface negative charges made it more difficult for carboxyl-QDs to adsorb on cell membrane.²² In addition, the result also indicated the greater uptake of *P. chrvsosporium* when treated ith smaller QDs (NH₂ CdSe/ZnS 525 and COOH CdSe/ZnS 525). Because the smaller CdSe/ZnS QDs were easier to be taken in by *P. chrysosporium*.⁴² Dae he small size, CdSe/ZnS QDs can directly enter into the hyphae by several approaches such as macropinocytosis, caveolae-mediated endocytosis, and elathrin-mediated endocytosis.⁴³ Meanwhile, the endocytosis is an energy consuming process, smaller CdSe/ZnS QDs would consume less energy than larger CoSe/ZnS QDs, leading to more uptake.⁴⁴ As shown in Fig. 2b, rere also observed by determining Se concentration, which the same results the accuracy of the employed ICP-OES measurement. Fig. 2c-d showed confirmed that there was good linear correlation between CdSe/ZnS QDs uptake and incubation concentration in the medium, indicating that the cellular uptake was a dose-dependent process.



CdSe/ZnS 525, and NH₂ CdSe/ZnS 625 by *A chrysosporium* incubated with different concentrations for 24 h. (a) CdSe/ZnS QDs, calculated from Cd, (b) CdSe/ZnS QDs, calculated from Se. Correlation between QDs uptake and incubation concentration in the medium. (c) CdSe/ZnS QDs, calculated from Cd, (d) CdSe/ZnS QDs, calculated from Se. Error bars represent one standard deviation of the arithmetic mean.

3.2.2. Incubation time

The incubation time plays a vital role in the uptake of QDs in *P. chrysosporium*. The cellular uptake of *P. chrysosporium* exposed to COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625 during 24 h was depicted in Fig. 3. As shown in Fig. 3a, the uptake of *P. chrysosporium* increased with

incubation time for all four types of CdSe/ZnS QDs. Since the accumulation of CdSe/ZnS QDs in P. chrysosporium presented a time-dependent process. However, the cellular uptake of different CdSe/ZnS QDs was obviously different under the identical conditions. For example, the cellular uptake of NH₂ CdSe/ZnS 525 reached 299.7 pmol g⁻¹ biomass at 24 h when the incubation concentration was 80 nM, which was greater than those of NH₂ CdSe/ZnS 625 (171.2 pmol g⁻¹ biomass), COOH CdSe/ZnS 525 (105.6 pmol g⁻¹ biomass), and COOH CdSe/ZnS 625 (72.8 phol g⁻¹ biomass), and the increase rate of NH₂ CdSe/ZnS 525 was largest. The cellular uptake of CdSe/ZnS QDs presented a time-dependent saturation, that was to say that the uptake amount of CdSe/ZnS QDs finally achieved a plateau. Because the cellular uptake process of CdSe/ZnS QDs included the briding of CdSe/ZnS QDs to receptors on the plasma membrane surface and the generation of coated pits to deliver them to the intracellular area.⁴⁵ The amount of receptors on plasma membrane surface would determine the uptake of GdSecond QDs, and the membrane receptors were gradually 45 Therefore, the internalization rate of CdSe/ZnS QDs consumed as time hosed mentually reach equilibrium. In addition, the smaller size and less would decline and negative barges make it easier and faster for NH2 CdSe/ZnS 525 to be internalized by *P. chrysosporium.*⁴⁴ As shown in Fig. 3b, the same results were also observed by determining Se concentration.



Fig. 3. Cellular uptake of COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625 by *P. chrysosporium* incubated for different

time. Initial CdSe/ZnS QDs concentrations are 10, 20, 50, and 80 nM, respectively. (a) CdSe/ZnS QDs, calculated from Cd, (b) CdSe/ZnS QDs, calculated from Se. Error bars represent one standard deviation of the arithmetic mean.

3.3. Confocal laser scanning microscopy analysis

To investigate the intracellular distribution of CdSe/ZnS QDs, the localization of CdSe/ZnS QDs in hyphae of P. chrysosporium were analyzed by confocal laser scanning microscopy. As shown in Fig. 4, the green fluorescence channels represented the hyphae of P. chrysosporium, in which all four types CdSe/ZnS QDs were largely accumulated. When CdSe/ZnS QDs entered to phae and accumulated largely in the hyphae of *P. chrysosporium*, RC absequently generated in vivo. as Therefore, the generation of intracellula qualitatively analyzed by confocal laser scanning microscopy coupled with H₂DCF-DA assay to explore the potential roles of CdSe/ZnS QDs in inducing intracellular oxidative stress. It can be seen from offluorescence indicated the generation of ROS in the Fig. 4 that the exi ium, and all samples presented significant differences as hyphae of P. ol. These phenomena suggested that CdSe/ZnS QDs accumulated compare to largely in the hyphae of P. chrysosporium, and the accumulated CdSe/ZnS QDs had induced the generation of ROS and caused oxidative stress in P. chrysosporium. Oxidative stress is one of the significant mechanisms of QDs cytotoxicity.^{46,47} The intracellular ROS would disturb the redox potential equilibrium, leading to an intracellular pro-oxidant environment, and ultimately cause the disruption of cell



CdSe/ZnS QDs exposure. Scale bars are 10 µm.

3.4.1. Lipid peroxidation and superoxide (O_2^{-}) production

MDA determination is generally used as an indicator of free radical production and a cytotoxic product of lipid peroxidation.^{35,49} In order to evaluate the CdSe/ZnS QDs-induced oxidative stress in *P. chrysosporium*, MDA content and O_2^- level in *P.* chrysosporium exposed to four types of CdSe/ZnS QDs were measured under different incubation concentrations. As shown in Fig. 5a, the MDA ontent in P. chrysosporium elevated obviously with incubation concentrations for all four types of CdSe/ZnS QDs. And the MDA content presented significant increase as compared to control under various concentrations. However the MDA contents of different CdSe/ZnS QDs exposure were distinctly different under the identical concentration. For example, the MDA content of MH_2 CdSe/ZnS 525 exposure reached 8.521 × 10⁻⁶ mol g⁻¹ when the incubation concentration was 80 nM, which was greater than those of NH₂ CdSe/ZnS 625 (6.95) × 10⁻⁶ mol g⁻¹), COOH CdSe/ZnS 525 (5.723 × 10⁻⁶ mol 625 (4.472 × 10⁻⁶ mol g⁻¹), and control (0.856 × 10⁻⁶ mol g⁻¹). g⁻¹), COOH CdSe/ZnS be related to the greatest cellular uptake of NH_2 CdSe/ZnS 525. The The result may increased MDA content showed that CdSe/ZnS QDs have stimulated the generation of free radical in *P. chrysosporium*. When CdSe/ZnS QDs entered into *P. chrysosporium*, they could inhibit the mitochondrial electron-transfer chain, resulting in the accumulation of semi-ubiquinone, which enabled to transfer one electron to molecular oxygen (O_2) to form superoxide radical (O_2) .⁵⁰ The generation of free radical, in turn,

led to lipid peroxidation.⁵¹ Thus, MDA content is associated with the production of O_2^- . As shown in Fig. 5b, the O_2^- level in *P. chrysosporium* exposed to four types of CdSe/ZnS QDs was similar to that of MDA, as described above. When *P. chrysosporium* was incubated with 80 nM CdSe/ZnS QDs for 24 h, the O_2^- level of NH₂ CdSe/ZnS 525 exposure reached 13.635 × 10⁻⁶ mol g⁻¹, which was greater than those of NH₂ CdSe/ZnS 625 (10.152 × 10⁻⁶ mol g⁻¹), COOH CdSe/ZnS 525 (8.823 × 10⁻⁶ mol g⁻¹), COOH CdSe/ZnS 625 (7.042 × 10⁻⁶ mol g⁻¹), and control (3213 × 10⁻⁶ mol g⁻¹). The increased O_2^- level is known as a significant factor for causing the cytotoxicity of CdSe/ZnS QDs. Since O_2^- has been proved to cause metabolic functions loss, DNA nicking and break, and apoptosis ¹⁵ herefore, the above results indicated that all four types of CdSe/ZnS QDs showed cytotoxicity to *P. chrysosporium* in the tested concentration and (10 - 80 nM).

3.4.2. Antioxidant enzymes analysis

Antioxidant enzymes have been considered as the defender in response to oxidative stress is in this study, two antioxidant enzymes of SOD and CAT have been monitored under different CdSe/ZnS QDs concentrations. As shown in Fig. 5c, the activity of SOD in *P. chrysosporium* increased gradually with incubation concentrations for all four types of CdSe/ZnS QDs. And the SOD activity presented significant increase as compared to control under various concentrations. However, the SOD activities of different CdSe/ZnS QDs exposure were distinctly different under the identical concentration. For example, the SOD activity of NH₂ CdSe/ZnS

525 exposure reached 9.235 U g^{-1} when the incubation concentration was 80 nM, which was greater than those of NH₂ CdSe/ZnS 625 (7.652 U g⁻¹), COOH CdSe/ZnS 525 (6.323 U g⁻¹), COOH CdSe/ZnS 625 (5.862 U g⁻¹), and control (2.304 U g⁻¹). The SOD activity decreased as follows: NH₂ CdSe/ZnS 525 > NH₂ CdSe/ZnS 625 > COOH CdSe/ZnS 525 > COOH CdSe/ZnS 625 > control, which suggested that NH₂ CdSe/ZnS 525 could induce the greater oxidative stress in P. chrysosporium than those of other three types of CdSe/ZnS QDs. Similarly, the result may ascribe to the greater cellular uptake of NH₂ CdSe/ZnS 525. Since the large amount of NH₂ CdSe/ZnS 525 in *P. chrysosporium* would lead to the accumulation of ROS.³⁰ The increased SOD activity was the physiological response of P. chrysosporium to As shown in Fig. 5d, the activity of CAT eliminate ROS and relieve oxidative stress. in P. chrysosporium exposed to four types of CdSe/ZnS QDs was similar to that of SOD, as described above. When R chrysosporium was incubated with 80 nM CdSe/ZnS QDs for 24 h the CAP activity of NH₂ CdSe/ZnS 525 exposure reached 49.250 U g⁻¹, which was greater than those of NH₂ CdSe/ZnS 625 (44.682 U g⁻¹), **52**5 (40.363 U g⁻¹), COOH CdSe/ZnS 625 (36.964 U g⁻¹), and COOH CdSe ZnS control (15.324 U g⁻¹). Therefore, the induction of antioxidant enzymes (SOD and CAT) activity is essential for P. chrysosporium to obtain the ability for relieving oxidative stress to some extent. Peng et al.⁴⁷ have reported that the antioxidant enzyme production might be a defense mechanism, since it provided a powerful defense against CdSe/ZnS QDs cytotoxicity before the induction of metallothionein synthesis.



Fig. 5. Oxidative stress in *P. chrysosporium* after exposure to COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625 with different concentrations for 24 h. The control group was the *P. chrysosporium* pellets without CdSe/ZnS QDs exposure (a) 4DA content, (b) O_2^- level, (c) SOD activity, (d) CAT activity. Error bars represent one standard deviation of the arithmetic mean.

3.4.3. Linear analysis of cytotoxicity response and CdSe/ZnS QDs concentrations

To explore the effect of CdSe/ZnS QDs exposure on *P. chrysosporium*, a linear analysis was used to reveal the relationship between cytotoxicity response of *P. chrysosporium* and CdSe/ZnS QDs concentrations. As depicted in Fig. 6a, MDA content, O_2^- level, SOD activity and CAT activity increased gradually with an increase of COOH CdSe/ZnS 525 concentrations, indicating that MDA content, O_2^- level, SOD

activity and CAT activity were dose-dependent and the cellular uptake of COOH CdSe/ZnS 525 increased with incubation concentrations. The similar results can be observed in Fig. 6b-d. However, there are also some differences among the four types of CdSe/ZnS QDs. For example, MDA content and O_2^- level were well correlated (R^2 = 0.98943 and R^2 = 0.97691, respectively) with the concentrations of COOH CdSe/ZnS 525 except for SOD activity and CAT activity ($R^2 = 0.91281$ and $R^2 =$ 0.89337, respectively). MDA content, O_2^- level and SOD activity were well correlated $(R^2 = 0.99557, R^2 = 0.97340 \text{ and } R^2 = 0.95507, \text{ respectively})$ with the concentrations of COOH CdSe/ZnS 625 except for CAT activity ($R^2 = 0.01532$) MDA content and O_2^- level were well correlated ($R^2 = 0.95270$ and $R^2 = 0.9973$, respectively) with the concentrations of NH₂ CdSe/ZnS 525 except for SOD activity and CAT activity ($R^2 =$ 0.79467 and $R^2 = 0.70510$, respectively). NOA content, O_2^- level and SOD activity were well correlated ($R^2 = 0.98238$, $R^2 = 0.99049$ and $R^2 = 0.95116$, respectively) with the concentrations of NH CdS47nS 625 except for CAT activity ($R^2 = 0.71448$). The antioxidant enzymes SOD and CAT) are produced to protect the cellular components from damage under CdSe/ZnS QDs-induced oxidative stress. However, when the induced xidetive stress exceed the scavenging ability of antioxidant enzymes, cellular damage would occur, confirmed by the decreased correlation between SOD and CAT activity and CdSe/ZnS QDs concentrations. These phenomena are in accordance with the findings reported by Chen et al.³⁰ who described the stress responses of P. chrysosporium under cadmium exposure and considered the slowly increased part as an exhaustion stage in which the defense systems were overloaded, resulting in chronic damage and cell death.



Fig. 6. Correlation between cytotoxicity response (MDA content, O_2^- level, SOD activity) and CAT activity) and CdSe/ZnS QDs concentration in the medium. (a) COOH CdSe/ZnS 525, (b) COOH CdSe/ZnS 625, (c) NH₂ CdSe/ZnS 525, (d) NH₂ CdSe/ZnS 625. R^2 represent the linearly dependent coefficient. Error bars represent one standard deviation of the arithmetic mean.

GSH is a significant antioxidant and can scavenge oxygen free radicals, and it plays an important role in the detoxification of exogenous chemicals.²² In order to investigate the CdSe/ZnS QDs-induced oxidative stress in P. chrysosporium, GSH content in P. chrysosporium exposed to four types of CdSe/ZnS QDs was measured under different incubation concentrations. As shown in Fig. 7, within the concentration range of 10 - 80 nM, there was an obvious reduction of CSH content in *P. chrysosporium* after incubation with all four types of CdSe/ZnS QDs. And the GSH content decreased significantly as compared to control undervarious concentrations. However, when P. chrysosporium was incubated with 80 rM CdSe/ZnS QDs for 24 h, the GSH content of NH₂ CdSe/ZnS 525 exposure was 41.5% of control, which was lower than those of NH₂ CdSe/ZnS 625 (\$6.2% of control), COOH CdSe/ZnS 525 Cise nS 625 (75.6% of control). The result showed (63.3% of control), and COOH that NH₂ CdSe/ZnS 525 has induced the greater oxidative stress in *P. chrysosporium* ypes of CdSe/ZnS QDs. The decreased GSH content most than those of othe probably attributed to the depletion of GSH in the scavenging of ROS,⁵⁵ which indicated the importance of GSH in detoxification and the ability of P. chrysosporium to tolerate CdSe/ZnS QDs exposure. The results are consistent with the report of Peng et al.²² who demonstrated direct evidence for the involvement of GSH in CdSe/ZnS QDs detoxification in HepG2 cells.



Fig. 7. The variation of GSH content in *P. chrysosporium* after exposure to COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625 with different concentrations for 24 h. The control group was the *P. chrysosporium* pellets without CdSe/ZnS QDs exposure. Error bars represent one standard deviation of the arithmetic mean.

4. Conclusions

This work comprehensively investigated the cellular uptake and induced physiological responses of four CdSe/ZnS QDs in *P. chrysosporium* via using ICP-OES, confocal laser scanning microscopy, and the analyses of MDA content, O_2^- level, SQD activity, CAT activity and GSH level. According to the measurement of ICP-OES, we found that negative charged CdSe/ZnS QDs with small size were more easily taken in by *P. chrysosporium* than large ones. Meanwhile, four CdSe/ZnS QDs accumulated largely in the hyphae and caused oxidative stress to *P. chrysosporium* in the tested concentration range. MDA content, O_2^- level, SOD activity, CAT activity and GSH level analysis results showed that CdSe/ZnS QDs with small size were more

cytotoxic than CdSe/ZnS QDs with large size, and small negative charged CdSe/ZnS QDs resulted in greater cytotoxicity than large negative charged CdSe/ZnS QDs. Further studies are urgently necessary to elucidate the underlying toxicity mechanism, which will eventually be used to prevent the adverse impacts of QDs in clinical application.

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgments

This work was financially supported by the National Natural Science Foundation of China (51521006, 51579099, 51508186 and 51378190), the Program for Changjiang Scholars and Innovative Research Team in University (IRT-13R17) and Hunan Provincial Natural Science Foundation of China (2015JJ2031 and 2016JJ3076).

References

- 1. C. Chan, D. J. Maxwell, X. Gao, R. E. Bailey, M. Han and S. Nie, *Curr. Opin. Biotech.*, 2002, **13**, 40-46.
- M. Chu, X. Pan, D. Zhang, Q. Wu, J. Peng and W. Hai, *Biomaterials*, 2012, 33, 7071-7083.
- 3. Y. Zhang, G. M. Zeng, L. Tang, D. L. Huang, X. Y. Jiang and Y. N. Chen,

Biosens. Bioelectron., 2007, 22, 2121-2126.

- X. Michalet, F. Pinaud, L. Bentolila, J. Tsay, S. Doose, J. Li, G. Sundaresan, A. Wu, S. Gambhir and S. Weiss, *Science*, 2005, **307**, 538-544.
- 5. I. L. Medintz, H. T. Uyeda, E. R. Goldman and H. Mattoussi, *Nat. Mater.*, 2005, **4**, 435-446.
- H. Wang, X. Z. Yuan, Y. Wu, X. H. Chen, L. J. Leng and G. M. Zeng, *RSC Adv.*, 2015, 5, 32531-32535.
- H. Wang, X. Z. Yuan, H. Wang, X. H. Chen, Z. B. Wu, L. B. Jiang, W. P. Xiong and G. M. Zeng, *Appl. Catal. B: Environ.*, 2016, 193, 36-46.
- Z. Z. Wu, X. Z. Yuan, H. Wang, Z. B. Wu, L. B. Jhang, H. Wang, L. Zhang, Z. H. Xiao, X. H. Chen and G. M. Zeng, *Appl. Catal. B: Environ.*, 2017, 202, 104-111.
- L. Hu, C. Zhang, G. Zeng, G. Chen, J. Wan, Z. Guo, H. Wu, Z. Yu, Y. Zhou and J. Liu, *RSC Adv.*, 2016, 6 78595-78610.
- 10. D. Mo, L. Hu, G. Zeng, G. Chen, J. Wan, Z. Yu, Z. Huang, K. He, C. Zhang and M. Cheng, *Appl. Microbiol. Biot.*, 2017, **101**, 2713-2733.
- L. Tang, G. M. Zeng, G. L. Shen, Y. P. Li, Y. Zhang and D. L. Huang, *Environ. Sci. Technol.*, 2008, 42, 1207-1212.
- M. D. Peterson, S. C. Jensen, D. J. Weinberg and E. A. Weiss, ACS Nano, 2014, 8, 2826-2837.
- N. A. Al-Hajaj, A. Moquin, K. D. Neibert, G. M. Soliman, F. O. M. Winnik and D. Maysinger, *Acs Nano*, 2011, 5, 4909-4918.

- Y. H. Luo, S. B. Wu, Y. H. Wei, Y. C. Chen, M. H. Tsai, C. C. Ho, S. Y. Lin, C.
 S. Yang and P. Lin, *Chem. Res. Toxicol.*, 2013, 26, 662-673.
- A. Nagy, A. Steinbru ck, J. Gao, N. Doggett, J. A. Hollingsworth and R. Iyer, Acs Nano, 2012, 6, 4748-4762.
- 16. G. Zeng, M. Chen and Z. Zeng, *Nature*, 2013, **499**, 154-154.
- P. Rivera-Gil, D. Jimenez De Aberasturi, V. Wulf, B. Pelaz, P. Del Pino, Y. Zhao, J. M. De La Fuente, I. Ruiz De Larramendi, T. Rojo and X J Liang, *Accounts Chem. Res.*, 2012, 46, 743-749.
- H. Li, M. Li, W. Y. Shih, P. I. Lelkes and W. H. Shih, J. Nanosci. Nanotechn., 2011, 11, 3543-3551.
- Y. Su, Y. He, H. Lu, L. Sai, Q. Li, W. D. Wang, P. Shen, Q. Huang and C. Fan, *Biomaterials*, 2009, **30**, 19-25
- 20. R. F. Domingos, D. F. Simon, C. Hauser and K. J. Wilkinson, *Environ. Sci. Technol.*, 2011, **45**, 7604-7669.
- S. J. Soenen, J. Demeester, S. C. De Smedt and K. Braeckmans, *Biomaterials*, 2012, 53, 4382-4888.
- 22. L. Peng, M. He, B. Chen, Q. Wu, Z. Zhang, D. Pang, Y. Zhu and B. Hu, *Biomaterials*, 2013, **34**, 9545-9558.
- 23. A. Hoshino, S. Hanada and K. Yamamoto, Arch. Toxicol., 2011, 85, 707.
- P. Xu, G. M. Zeng, D. L. Huang, C. L. Feng, S. Hu, M. H. Zhao, C. Lai, Z. Wei,
 C. Huang and G. X. Xie, *Sci. Total Environ.*, 2012, **424**, 1-10.
- 25. C. Yang, H. Chen, G. Zeng, G. Yu and S. Luo, Biotechnol. Adv., 2010, 28,

531-540.

- T. Fan, Y. Liu, B. Feng, G. Zeng, C. Yang, M. Zhou, H. Zhou, Z. Tan and X. Wang, *J. Hazard. Mater.*, 2008, 160, 655-661.
- D. L. Huang, G. M. Zeng, C. L. Feng, S. Hu, X. Y. Jiang, L. Tang, F. F. Su, Y. Zhang, W. Zeng and H. L. Liu, *Environ. Sci. Technol.*, 2008, 42, 4946-4951.
- 28. Y. Feng, J. L. Gong, G. M. Zeng, Q. Y. Niu, H. Y. Zhang, C. G. Niu, J. H. Deng and M. Yan, *Chem. Eng. J.*, 2010, 162, 487-494.
- 29. G. Zeng, M. Chen and Z. Zeng, Science, 2013, 340, 1403.
- 30. A. Chen, G. Zeng, G. Chen, L. Liu, C. Shang, X. Ha, L. Lu, M. Chen, Y. Zhou and Q. Zhang, *Process Biochem.*, 2014, 49, 583-598
- 31. L. Hu, G. Zeng, G. Chen, H. Dong, Y Du J. Wan, A. Chen, Z. Guo, M. Yan and H. Wu, *J. Hazard. Mater.*, 2016 **301**, 106-118.
- 32. J. L. Gong, B. Wang, G. M. Zeng, C. P. Yang, C. G. Niu, Q. Y. Niu, W. J. Zhou and Y. Liang, J. Hagara Mater., 2009, 164, 1517-1522.
- Y. He, H. T. Iu, L. M. Sai, Y. Y. Su, M. Hu, C. H. Fan, W. Huang and L. H.
 Wang Adv. Mater., 2008, 20, 3416-3421.
- 34. T.K. Birk, E. Schultz, W. Connors, L. Lorenz and J. Zeikus, *Arch. Microbiol.*, 1978, 117, 277-285.
- 35. G. M. Zeng, A. W. Chen, G. Q. Chen, X. J. Hu, S. Guan, C. Shang, L. H. Lu and Z. J. Zou, *Environ. Sci. Technol.*, 2012, 46, 7818-7825.
- 36. P. Aravind and M. N. V. Prasad, *Plant Physiol. Bioch.*, 2003, **41**, 391-397.
- 37. Y. Lei, C. Yin and C. Li, *Physiol. Plantarum*, 2006, **127**, 182-191.

- F. R. Cavalcanti, J. T. A. Oliveira, A. S. Martins-Miranda, R. A. Viégas and J.
 A. G. Silveira, *New Phytol.*, 2004, 163, 563-571.
- 39. A. Rehman and M. S. Anjum, Environ. Monit. Asses., 2011, 174, 585-595.
- 40. Y. Su, F. Peng, Z. Jiang, Y. Zhong, Y. Lu, X. Jiang, Q. Huang, C. Fan, S. T. Lee and Y. He, *Biomaterials*, 2011, **32**, 5855-5862.
- 41. X. J. Hu, J. S. Wang, Y. G. Liu, X. Li, G. M. Zeng, Z. L. Bao, X. X. Zeng, A. W.
 Chen and F. Long, *J. Hazard. Mater.*, 2011, 185, 306-314.
- 42. Z. Huang, G. Chen, G. Zeng, Z. Guo, K. He, L. Hu, J. Wu, L. Zhang, Y. Zhu and Z. Song, J. Hazard. Mater., 2017, **321**, 37-46.
- S. Zhang, Y. Jiang, C. S. Chen, J. Spurgin, K. A. Schwehr, A. Quigg, W. C. Chin and P. H. Santschi, *Environ. Sci. Technol.*, 2012, 46, 8764-8772.
- 44. X. Jiang, C. Rocker, M. Hafner, S. Brandholt, R. M. Dorlich and G. U. Nienhaus, ACS Nano, 2010, 4, 6787-6797.
- 45. S. Ohta, S. Inasawa and Yamaguchi, *Biomaterials*, 2012, **33**, 4639-4645.
- E. Q. Contrens, M. Cho, H. Zhu, H. L. Puppala, G. Escalera, W. Zhong and V.
 L. Colvin, *Environ. Sci. Technol.*, 2013, 47, 1148.
- 47. L. Perg, M. He, B. Chen, Y. Qiao and B. Hu, ACS Nano, 2015, 9, 10324-10334.
- J. Wan, G. Zeng, D. Huang, C. Huang, C. Lai, N. Li, Z. Wei, P. Xu, X. He and M. Lai, *Appl. Biochem. Biotech.*, 2015, **175**, 1981-1991.
- J. Wan, C. Zhang, G. Zeng, D. Huang, L. Hu, C. Huang, H. Wu and L. Wang, J. Hazard. Mater., 2016, 320, 278-288.

- S. Garg, H. Rong, C. J. Miller and T. D. Waite, *Environ. Sci. Technol.*, 2016, 50, 3890-3896.
- Y. Lu, S. Xu, H. Chen, M. He, Y. Deng, Z. Cao, H. Pi, C. Chen, M. Li and Q. Ma, *Biomaterials*, 2016, **90**, 27-39.
- 52. R. Liu, H. H. Liu, Z. Ji, C. H. Chang, T. Xia, A. E. Nel and Y. Cohen, ACS Nano, 2015, **9**, 9303-9313.
- Y. Cheng, H. He, C. Yang, G. Zeng, X. Li, H. Chen and G. Yu, *Biotechnol. Adv.*, 2016, 34, 1091-1102.
- G. M. Zeng, J. Wan, D. L. Huang, L. Hu, C. Huang, M. Cheng, W. J. Xue, X.
 M. Gong, R. Z. Wang and D. N. Jiang, *J. Hazard. Mater.*, 2017, 339, 354-367.
- 55. Y. Matsufuji, K. Yamamoto, K. Yamanon, N. Mitsunaga, T. Hayakawa and T. Nakagawa, *Appl. Microbiol. Biotectural.*, 2013, **97**, 297-303.

List of figures

Fig. 1. High-resolution TEM mages of COOH CdSe/ZnS 525 (a), COOH CdSe/ZnS 625 (b), NH₂ CdSe/ZrS 515 (c), and NH₂ CdSe/ZnS 625 (d), respectively.

Fig. 2. Cellular hptake of COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/Zn, 523, and NH₂ CdSe/ZnS 625 by *P. chrysosporium* incubated with different concentrations for 24 h. (a) CdSe/ZnS QDs, calculated from Cd, (b) CdSe/ZnS QDs, calculated from Se. Correlation between QDs uptake and incubation concentration in the medium. (c) CdSe/ZnS QDs, calculated from Cd, (d) CdSe/ZnS QDs, calculated from Se. Error bars represent one standard deviation of the arithmetic mean.

Fig. 3. Cellular uptake of COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂

CdSe/ZnS 525, and NH₂ CdSe/ZnS 625 by *P. chrysosporium* incubated for different time. Initial CdSe/ZnS QDs concentrations are 10, 20, 50, and 80 nM, respectively. (a) CdSe/ZnS QDs, calculated from Cd, (b) CdSe/ZnS QDs, calculated from Se. Error bars represent one standard deviation of the arithmetic mean.

Fig. 4. Confocal laser scanning microscopy images of the hyphal localization (green fluorescence channel) of COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625. And their initial concentrations are 10/20, 50, and 80 nM, respectively. The control group was the *P. chrysosporhum* pellets without CdSe/ZnS QDs exposure. Scale bars are 10 μm.

Fig. 5. Oxidative stress in *P. chrysosporium* after exposure to COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625 with different concentrations for 24 h. The control group was the *P. chrysosporium* pellets without CdSe/ZnS QDs exposure. (a) MIIA content, (b) O_2^- level, (c) SOD activity, (d) CAT activity. Error bars represent one standard deviation of the arithmetic mean.

Fig. 6. Correlation between cytotoxicity response (MDA content, O_2^- level, SOD activity, and CAT activity) and CdSe/ZnS QDs concentration in the medium. (a) COOH CdSe/ZnS 525, (b) COOH CdSe/ZnS 625, (c) NH₂ CdSe/ZnS 525, (d) NH₂ CdSe/ZnS 625. R^2 represent the linearly dependent coefficient. Error bars represent one standard deviation of the arithmetic mean.

Fig. 7. The variation of GSH content in *P. chrysosporium* after exposure to COOH CdSe/ZnS 525, COOH CdSe/ZnS 625, NH₂ CdSe/ZnS 525, and NH₂ CdSe/ZnS 625 with different concentrations for 24 h. The control group was the *P. chrysosporium*

pellets without CdSe/ZnS QDs exposure. Error bars represent one standard deviation of the arithmetic mean.



Table of Contents